arXiv:0904.2871v1 [math.SP] 18 Apr 2009

1D SCHRÖDINGER OPERATOR WITH PERIODIC PLUS COMPACTLY SUPPORTED POTENTIALS

EVGENY KOROTYAEV

Abstract. We consider the 1D Schrödinger operator $Hy = -y'' + (p+q)y$ with a periodic potential p plus compactly supported potential q on the real line. The spectrum of H consists of an absolutely continuous part plus a finite number of simple eigenvalues in each spectral gap $\gamma_n \neq \emptyset, n \geqslant 0$, where γ_0 is unbounded gap. We prove the following results: 1) we determine the distribution of resonances in the disk with large radius, 2) a forbidden domain for the resonances is specified, 3) the asymptotics of eigenvalues and antibound states are determined, 4) if $q_0 = \int_{\mathbb{R}} q dx = 0$, then roughly speaking in each nondegenerate gap γ_n for n large enough there are two eigenvalues and zero antibound state or zero eigenvalues and two antibound states, 5) if H has infinitely many gaps in the continuous spectrum, then for any sequence $\sigma = (\sigma)_1^\infty, \sigma_n \in \{0,2\}$, there exists a compactly supported potential q such that H has σ_n bound states and $2-\sigma_n$ antibound states in each gap γ_n for n large enough. 6) For any q (with $q_0 = 0$), $\sigma = (\sigma_n)_1^\infty$, where $\sigma_n \in \{0,2\}$ and for any sequence $\delta = (\delta_n)_1^\infty \in \ell^2, \delta_n > 0$ there exists a potential $p \in L^2(0,1)$ such that each gap length $|\gamma_n| = \delta_n, n \geqslant 1$ and H has exactly σ_n eigenvalues and $2-\sigma_n$ antibound states in each gap $\gamma_n \neq \emptyset$ for n large enough.

1. Introduction

Consider the Schrödinger operator $H = H_0 + q$ acting in $L^2(\mathbb{R})$, where $H_0 = -\frac{d^2}{dx^2} + p(x)$ and $p \in L^2(0,1)$ is the real 1-periodic potential. The real compactly supported potential q belongs to the class $\mathcal{Q}_t = \{q \in L^2(\mathbb{R}) : \ [0,t] \ \textit{is the convex hull of the support of } q\}$ for some $t > 0$. The spectrum of H_0 consists of spectral bands $\mathfrak{S}_n$ and is given by (see Fig. 1)

$$\sigma(H_0) = \sigma_{ac}(H_0) = \cup \mathfrak{S}_n, \quad \mathfrak{S}_n = [E_{n-1}^+, E_n^-], n \geqslant 1, \quad E_0^+ < .. \leqslant E_{n-1}^+ < E_n^- \leqslant E_n^+ < ...$$

We assume that $E_0^+ = 0$. The sequence $E_0^+ < E_1^- \leqslant E_1^+ < \dots$ is the spectrum of the equation

$$-y'' + p(x)y = \lambda y, \qquad \lambda \in \mathbb{C}, \tag{1.1}$$

with the 2-periodic boundary conditions, i.e. $y(x+2) = y(x), x \in \mathbb{R}$. The bands $\mathfrak{S}_n, \mathfrak{S}_{n+1}$ are separated by a gap $\gamma_n = (E_n^-, E_n^+)$ and let $\gamma_0 = (-\infty, E_0^+)$. If a gap degenerates, that is $\gamma_n = \emptyset$, then the corresponding bands $\mathfrak{S}_n$ and $\mathfrak{S}_{n+1}$ merge. If $E_n^- = E_n^+$ for some n, then this number $E_n^\pm$ is the double eigenvalue of equation (1.1) with the 2-periodic boundary conditions. The lowest eigenvalue $E_0^+ = 0$ is always simple and the corresponding eigenfunction is 1-periodic. The eigenfunctions, corresponding to the eigenvalue $E_{2n}^\pm$, are 1-periodic, and for the case $E_{2n+1}^\pm$ they are anti-periodic, i.e., $y(x+1) = -y(x), \quad x \in \mathbb{R}$.

It is well known, that the spectrum of H consists of an absolutely continuous part $\sigma_{ac}(H) = \sigma_{ac}(H_0)$ plus a finite number of simple eigenvalues in each gap $\gamma_n \neq \emptyset, n \geqslant 0$, see [Rb], [F1]. Moreover, in every open gap γ_n for n large enough the operator H has at most two bound

---

Date: April 20, 2009.

1991 *Mathematics Subject Classification.* 34A55, (34B24, 47E05).

Key words and phrases. resonances, scattering, periodic potential, S-matrix.



FIGURE 1. The cut domain $\mathbb{C}\setminus\cup\mathfrak{S}_n$ and the cuts (bands) $\mathfrak{S}_n=[E_{n-1}^+,E_n^-], n\geqslant 1$

states [Rb] and precisely one bound state in the case $q_0=\int_{\mathbb{R}}q(x)dx\neq 0$ [Zh], [F2]. Note that the potential q in [Rb],[Zh] belongs to the more general class, see also [GS], [So].

Let $\varphi(x,z),\vartheta(x,z)$ be the solutions of the equation $-y''+py=z^2y$ satisfying $\varphi'(0,z)=\vartheta(0,z)=1$ and $\varphi(0,z)=\vartheta'(0,z)=0$, where $u'=\partial_x u$. The Lyapunov function is defined by $\Delta(z)=\frac{1}{2}(\varphi'(1,z)+\vartheta(1,z))$. The function $\Delta^2(\sqrt{\lambda})$ is entire, where $\sqrt{\lambda}$ is defined by $\sqrt{1}=1$. Introduce the function $\Omega(\lambda)=(1-\Delta^2(\sqrt{\lambda}))^{\frac{1}{2}}, \lambda\in\overline{\mathbb{C}}_+$ and we fix the branch $\Omega(\lambda)=(1-\Delta^2(\sqrt{\lambda}))^{\frac{1}{2}}$ by the condition $\Omega(\lambda+i0)>0$ for $\lambda\in\mathfrak{S}_1=[E_0^+,E_1^-]$. Introduce the two-sheeted Riemann surface Λ of $\Omega(\lambda)=(1-\Delta^2(\sqrt{\lambda}))^{\frac{1}{2}}$ obtained by joining the upper and lower rims of two copies of the cut plane $\mathbb{C}\setminus\sigma_{ac}(H_0)$ in the usual (crosswise) way. The n-th gap on the first physical sheet Λ_1 we will denote by $\gamma_n^{(1)}$ and the same gap but on the second nonphysical sheet Λ_2 we will denote by $\gamma_n^{(2)}$ and let $\gamma_n^{(0)}$ be the union of $\overline{\gamma}_n^{(1)}$ and $\overline{\gamma}_n^{(2)}$, i.e.,

$$\gamma_n^{(0)}=\overline{\gamma}_n^{(1)}\cup\overline{\gamma}_n^{(2)}.$$

Introduce the function $D(\lambda)=\det(I+q(H_0-\lambda)^{-1})$, which is meromorphic in Λ, see [F1]. Recall that $1/D$ is the transmission coefficient in the S-matrix for the operators H,H_0, see Sect. 2. If D has some poles, then they coincide with some $E_n^\pm$. It is well known that if $D(\lambda)=0$ for some zero $\lambda\in\Lambda_1$, then λ is an eigenvalue of H and $\lambda\in\cup\gamma_n^{(1)}$. Note that there are no eigenvalues on the spectrum $\sigma_{ac}(H_0)\subset\Lambda_1$, since $|D(\lambda)|\geqslant 1$ on $\sigma_{ac}(H_0)\subset\Lambda_1$ (see (2.19) and all these facts in [F1]).

Define the functions A,J by

$$J(\lambda)=2\Omega(\lambda+i0)\operatorname{Im}D(\lambda+i0),\qquad A(\lambda)=\operatorname{Re}D(\lambda+i0)-1,\quad for\quad \lambda\in\sigma(H_0)\subset\Lambda_1. \tag{1.2}$$

In Lemma 3.2 we will show that A,J are entire functions on $\mathbb{C}$ and they are real on the real line. Instead of the function D we will study the **modified function** $\Xi=2i\Omega D$ on Λ. We will show that Ξ satisfies

$$\Xi=2i\Omega D=2i\Omega(1+A)-J\qquad on\qquad \Lambda, \tag{1.3}$$

see Theorem 1.1. Recall that Ω is analytic in Λ and $\Omega(\lambda)=0$ for some $\lambda\in\Lambda$ iff $\lambda=E_n^-$ or $\lambda=E_n^+$ for some $n\geqslant 0$. Then the function Ξ is analytic on Λ and has branch points $E_n^\pm,\gamma_n\neq\emptyset$. The zeros of Ξ define the eigenvalues and resonances, similar to the case $p=0$. Define the set $\Lambda_0=\{\lambda\in\Lambda:\lambda=E_n^+\in\Lambda_1 \text{ and } \lambda=E_n^+\in\Lambda_2,\gamma_n=\emptyset,n\geqslant 1\}\subset\Lambda$. In fact with each $\gamma_n=\emptyset$ we associate two points $E_n^+\in\Lambda_1$ and $E_n^+\in\Lambda_2$ from the set Λ_0. If each gap of H_0 is empty, then Λ_0 is a union of two sets $\{(\pi n)^2,n\in\mathbb{N}\}\subset\Lambda_1$ and $\{(\pi n)^2,n\in\mathbb{N}\}\subset\Lambda_2$. If each gap of H_0 is not empty, then $\Lambda_0=\emptyset$.

Definition of states. *Each zero of* Ξ *in* $\Lambda\setminus\Lambda_0$ *is a state of* H.

1) A state $\lambda\in\Lambda_2$ *is a resonance.*

2) A state $\lambda = E_n^\pm, n \geqslant 0$ *is a virtual state.*
3) A resonance $\lambda \in \cup\gamma_n^{(2)} \subset \Lambda_2$ *is an anti-bound state.*

It is known that the gaps $\gamma_n = \emptyset$ do not give contribution to the states. Recall that S-matrix for H, H_0 is meromorphic on Λ, but it is analytic at the points from Λ_0 (see [F1]). Roughly speaking there is no difference between the points from Λ_0 and other points inside the spectrum of H_0.

The gap $\gamma_n^{(1)} \subset \Lambda_1$ is so-called physical gap and the gap $\gamma_n^{(2)} \subset \Lambda_2$ is so-called non-physical gap. If $q_0 = \int_{\mathbb{R}} q(x)dx \neq 0$, then H has precisely one bound state on each physical gap $\neq \emptyset$ and odd number $\geqslant 1$ of resonances on each non-physical gap $\neq \emptyset$ for n large enough [F1].

We explain roughly why antibound states are important. Consider the operator $H_\tau = H_0 + \tau q$, where $\tau \in \mathbb{R}$ is the coupling constant. If $\tau = 0$, then $H = H_0$ and there are no eigenvalues, complex resonances and H has only virtual states, which coincide with each $E_n^\pm, , \gamma_n \neq \emptyset$, since $\Xi = 2i\Omega$ on Λ at $q = 0$ and $\Omega(E_n^\pm) = 0$. If τ is increasing, then there are eigenvalues, antibound states (close to the end of gaps) and complex resonances. If τ is increasing again, then there are no new eigenvalues but some complex resonances ($\zeta \in \mathbb{C}_+ \subset \Lambda_2$ and $\overline{\zeta} \in \mathbb{C}_- \subset \Lambda_2$) reach some non physical gap and transform into new antibound states. If τ is increasing again, then some new antibound states will be virtual states, and then later they will be bound states. Thus if τ runs through $\mathbb{R}_+$, then there is a following transformation: resonances $\to$ antibound states $\to$ virtual states $\to$ bound states $\to$ virtual states...

For each $n \geqslant 1$ there exists an unique point $E_n \in [E_n^-, E_n^+]$ such

$$(-1)^n \Delta(\sqrt{E_n}) = \max_{z \in [E_n^-, E_n^+]} |\Delta(\sqrt{\lambda})| = \cosh h_n, \qquad \text{for some } h_n \geqslant 0. \tag{1.4}$$

Note that if $\gamma_n = \emptyset$, then $E_n = E_n^\pm$ and if $\gamma_n \neq \emptyset$, then $E_n \in \gamma_n$.

Theorem 1.1. *Let potentials* $(p, q) \in L^2(0,1) \times \mathcal{Q}_t, t > 0$. *Then* Ξ *satisfies (1.3) and*
i) There exist even number $\geqslant 0$ *of states (counted with multiplicity) on each set* $\gamma_n^{(0)} \neq \emptyset, n \geqslant 1$, *where* $\gamma_n^{(0)}$ *is a union of the physical* $\overline{\gamma}_n^{(1)} \subset \Lambda_1$ *and non-physical gap* $\overline{\gamma}_n^{(2)} \subset \Lambda_2$ *(here* $\gamma_n \neq \emptyset$*).*
ii) Let $\|q\|_t = \int_0^t |q(x)|dx$. *There are no states in the "forbidden" domain* $\mathscr{D}_F \setminus \cup\overline{\gamma}_n^{(2)} \subset \Lambda_2$, *where*

$$\mathscr{D}_F = \{\lambda \in \Lambda_2 : |\lambda|^{\frac{1}{2}} > \max\{180 e^{2\|p\|_1}, C_F e^{2t|\operatorname{Im}\sqrt{\lambda}|}\}\}, \quad C_F = 12\|q\|_t e^{\|p\|_1 + \|q\|_t + 2\|p\|_t}. \tag{1.5}$$

iii) In each $\gamma_n^{(0)} \neq \emptyset, n \geqslant 1 + \frac{e^{t\pi/2}}{\pi} C_F$ *there exists exactly two simple real states* $\lambda_n^\pm \in \gamma_n^{(0)}$ *such that* $E_n^- \leqslant \lambda_n^- < E_n < \lambda_n^+ \leqslant E_n^+$.
iv) Let $\lambda \in \gamma_n^{(1)}$ *be an eigenvalue of* H *for some* $n \geqslant 0$, *i.e.,* $\Xi(\lambda) = 0$ *and let* $\lambda^{(2)} \in \gamma_n^{(2)} \subset \Lambda_2$ *be the same number but on the second sheet* Λ_2. *Then* $\lambda^{(2)}$ *is not an anti-bound state, i.e.,* $\Xi(\lambda^{(2)}) \neq 0$.

Remark. 1) The forbidden domain $\mathscr{D}_F \cap \mathbb{C}_-$ is similar to the case $p = 0$, see [K2].
2) In Theorem 1.2 we show that $\lambda_n^\pm \to E_n^\pm$ as $n \to \infty$.

Let $\mu_n^2, n \geqslant 1$ be the Dirichlet spectrum of the equation $-y'' + py = \mu^2 y$ on the interval $[0,1]$ with the boundary condition $y(0) = y(1) = 0$. It is well known that each $\mu_n^2 \in [E_n^-, E_n^+], n \geqslant 1$. Define the coefficients $q_0 = \int_{\mathbb{R}} q(x)dx$ and

$$\widehat{q}_n = \widehat{q}_{cn} + i\widehat{q}_{sn}, \quad \widehat{q}_{cn} = \int_{\mathbb{R}} q(x)\cos 2\pi nx dx, \qquad \widehat{q}_{sn} = \int_{\mathbb{R}} q(x)\sin 2\pi nx dx, \ n \geqslant 1. \tag{1.6}$$

In order to formulate Theorem 1.2 we define the functions $\Psi_{cn}, \Psi_{sn}, c_n = \cos\phi_n, s_n = \sin\phi_n, n \geqslant 1$ (depending from $p \in L^2(0,1)$) by

$$\Psi_{cn} = \frac{E_n^- + E_n^+}{2} - \mu_n^2 = \frac{|\gamma_n|}{2}c_n, \quad \Psi_{sn} = \frac{|\gamma_n|}{2}s_n, \quad \operatorname{sign} s_n = \operatorname{sign}|\varphi'(1,\mu_n)|. \tag{1.7}$$

The identity $\frac{E_n^- + E_n^+}{2} - \mu_n^2 = \frac{|\gamma_n|}{2}c_n$ defines $c_n = \cos\phi_n \in [-1,1]$ and the identity $\operatorname{sign} s_n = \operatorname{sign}|\varphi'(1,\mu_n)|$ defines $s_n = \sin\phi_n$ and the angle $\phi_n \in [0, 2\pi)$. Recall the results from [K5]:

The mapping $\Psi : \mathcal{H} \to \ell^2 \oplus \ell^2$ *given by* $\Psi = ((\Psi_{cn})_1^\infty, (\Psi_{sn})_1^\infty)$ *is a real analytic isomorphism between real Hilbert spaces* $\mathcal{H} = \{p \in L^2(0,1) : \int_0^1 p(x)dx = 0\}$ *and* $\ell^2 \oplus \ell^2$.

Let $\#(H, r, X)$ be the total number of state of H in the set $X \subseteq \Lambda$ having modulus $\leqslant r^2$, each state being counted according to its multiplicity.

Theorem 1.2. *Let* $(p,q) \in L^2(0,1) \times \mathcal{Q}_t$ *and let* $q_0 = \int_{\mathbb{R}} q(x)dx$. *Then*
i) Let $\gamma_n \neq \emptyset$ *for some* n *large enough. Then* H *has exactly two simple states* $\lambda_n^-, \lambda_n^+ \in \gamma_n^{(0)}$. *Moreover, if* λ *is one of* λ_n^-, λ_n^+ *and satisfies* $(-1)^{n+1}J(\lambda) > 0$ *(or* $(-1)^{n+1}J(\lambda) < 0$ *or* $J(\lambda) = 0$*), then* λ *is a bound state (or an anti-bound state or a virtual state).*
ii) The following asymptotics hold true:

$$\sqrt{\lambda_n^\pm} = \sqrt{E_n^\pm} \mp \frac{2|\gamma_n|}{(4\pi n)^3}(\mp q_0 - c_n\widehat{q}_{cn} + s_n\widehat{q}_{sn} + O(1/n))^2,$$

$$(-1)^{n+1}J(\lambda_n^\pm) = \frac{|\gamma_n|}{(2\pi n)^2}(\mp q_0 - c_n\widehat{q}_{cn} + s_n\widehat{q}_{sn} + O(1/n)),$$

$$\begin{aligned} &\textit{if} \quad q_0 > 0 \ \Rightarrow \quad \lambda_n^- \in \Lambda_1 \textit{ is bound state}, \qquad \lambda_n^+ \in \Lambda_2 \textit{ is anti bound state},\\ &\textit{if} \quad q_0 < 0 \ \Rightarrow \quad \lambda_n^- \in \Lambda_2 \textit{ is anti bound state}, \qquad \lambda_n^+ \in \Lambda_1 \textit{ is bound state}. \end{aligned} \tag{1.8}$$

iii) Let $\widehat{q}_n = \widehat{q}_{cn} + i\widehat{q}_{sn} = |\widehat{q}_n|e^{i\tau_n}, n \geqslant 1$ *and* $q_0 = 0$. *Assume that* $|\cos(\phi_n + \tau_n)| > \varepsilon > 0$ *and* $|\widehat{q}_n| > n^{-\alpha}$ *for* n *large enough and for some* $\varepsilon, \alpha \in (0,1)$. *Then the operator* H *has* $\sigma_n = 1 - \operatorname{sign}\cos(\phi_n + \tau_n)$ *bound states in the physical gap* $\gamma_n^{(1)} \neq \emptyset$ *and* $2 - \sigma_n$ *resonances inside the nonphysical gap* $\gamma_n^{(2)} \neq \emptyset$ *for* n *large enough.*
iv) For some integer $N_S \in \mathbb{Z}$ *the following asymptotics hold true:*

$$\#(H, r, \Lambda_2 \setminus \cup\gamma_n^{(2)}) = r\frac{2t + o(1)}{\pi} \quad \textit{as} \quad r \to \infty, \tag{1.9}$$

$$\#(H, r, \cup\Gamma_n) = \#(H_0, r, \cup\Gamma_n) + 2N_S \quad \textit{as} \quad r \to \infty, \quad r^2 \notin \cup\overline{\gamma}_n. \tag{1.10}$$

Remark. 1) First term in the asymptotics (1.9) does not depend on the periodic potential p. Recall that asymptotics (1.9) for the case $p = 0$ was obtained by Zworski [Z].
2) The main difference between the distribution of the resonances for the case $p \neq$ const and $p =$ const is the bound states and anti bound states in high energy gaps, see (iii) and (1.8).
3) Assume that a potential $u \in L^2(\mathbb{R})$ is compactly supported, $\operatorname{supp} u \subset (0,t)$ and satisfies $|\widehat{u}_n| = o(n^{-\alpha})$ as $n \to \infty$. Then in the case (iii) the operator $H + u$ has also $1 - \operatorname{sign}\cos(\phi_n + \tau_n)$ bound states in each gap $g_n \neq \emptyset$ for n large enough.
4) In the proof of (1.9) we use the Paley Wiener type Theorem from [Fr], the Levinson Theorem (see Sect. 3) and a priori estimates from [KK], [M].

5) Assume that $\mu_n^2 = E_n^-$ or $\mu_n^2 = E_n^+$ with $|\gamma_n| > 0$ for all $n \in \mathbb{N}_0$, where $\mathbb{N}_0 \subset \mathbb{N}$ is some infinite subset. Then for all large $n \in \mathbb{N}_0$ the following asymptotics hold true:

$$\sqrt{\lambda_n^\pm} = \sqrt{E_n^\pm} \mp \frac{2|\gamma_n|}{(4\pi n)^2}\biggl(\mp q_0 - c_n \widehat{q}_{cn} + \frac{O(1)}{n}\biggr)^2 \qquad \text{as} \quad n \to \infty. \tag{1.11}$$

There is no $\widehat{q}_{sn}$ in asymptotics (1.11). Let in addition $q_0 = 0$ and $|\widehat{q}_{cn}| > n^{-\alpha}$ for all $n \in \mathbb{N}_0$ and some $\alpha \in (0,1)$. Then H has exactly $\sigma_n = 1 - \operatorname{sign} c_n \widehat{q}_{cn}$ eigenvalues in each gap $\gamma_n^{(1)}$ and $2 - \sigma_n$ resonances inside each gap $\gamma_n^{(2)} \neq \emptyset$ for $n \in \mathbb{N}_0$ large enough.
6) Recall that if p is even i.e., $p \in L^2_{even}(0,1) = \{p \in L^2(0,1), p(x) = p(1-x), x \in (0,1)\}$, iff $s_n = 0$ (or $\mu_n^2 \in \{E_n^-, E_n^+\}$, or $c_n \in \{\pm 1\}$) for all $n \geqslant 1$, see [GT], [KK1].

Consider some inverse problems for the operator H.

Theorem 1.3. *i) Let the operator H_0 have infinitely many gaps $\gamma_n \neq \emptyset$ for some $p \in L^2(0,1)$. Then for any sequence $\sigma = (\sigma_n)_1^\infty$, where $\sigma_n \in \{0,2\}$, there exists some potential $q \in \mathcal{Q}^t, t > 0$ such that H has exactly σ_n eigenvalues in each gap $\gamma_n^{(1)} \neq \emptyset$ and $2 - \sigma_n$ resonances inside each gap $\gamma_n^{(2)} \neq \emptyset$ for n large enough.*
ii) Let $q \in \mathcal{Q}_t, t > 0$ satisfy $q_0 = 0$ and let $|\widehat{q}_{cn}| > n^{-\alpha}$ for all n large enough and some $\alpha \in (0,1)$. Let $\sigma = (\sigma_n)_1^\infty$, where $\sigma_n \in \{0,2\}$ and infinitely many $\delta_n > 0$. Then for any sequence $\delta = (\delta_n)_1^\infty \in \ell^2$ where each $\delta_n \geqslant 0, n \geqslant 1$ there exists a potential $p \in L^2(0,1)$ such that each gap length $|\gamma_n| = \delta_n, n \geqslant 1$. Moreover, H has exactly σ_n eigenvalues in each physical gap $\gamma_n^{(1)} \neq \emptyset$ and $2 - \sigma_n$ resonances inside each non-physical gap $\gamma_n^{(2)} \neq \emptyset$ for n large enough.

Remark . 1) In the proof we use results from the inverse spectral theory from [K5].
2) We have $\alpha \in (\frac12, 1)$, if all gaps are open.

A lot of papers are devoted to the resonances for the Schrödinger operator with $p = 0$, see [Fr], [H], [K1], [K2], [S], [Z] and references therein. Although resonances have been studied in many settings, but there are relatively few cases where the asymptotics of the resonance counting function are known, mainly one dimensional case [Fr], [K1], [K2], [S], and [Z]. We recall that Zworski [Z] obtained the first results about the distribution of resonances for the Schrödinger operator with compactly supported potentials on the real line. The author obtained the characterization (plus uniqueness and recovering) of S-matrix for the Schrödinger operator with a compactly supported potential on the real line [K2] and the half-line [K1], see also [Z1], [BKW] about uniqueness.

For the Schrödinger operator on the half line the author [K3] obtained the stability results:

(i) If $\varkappa^0 = (\varkappa^0)_1^\infty$ is a sequence of zeros (eigenvalues and resonances) of the Jost function for some real compactly supported potential q_0 and $\varkappa - \varkappa^0 \in \ell^2_\varepsilon$ for some $\varepsilon > 1$, then $\varkappa$ is the sequence of zeroes of the Jost function for some unique real compactly supported potential.

(ii) The measure associated with the zeros of the Jost function is the Carleson measure, the sum $\sum (1 + |\varkappa_n^0|)^{-\alpha}, \alpha > 1$ is estimated in terms of the L^1-norm of the potential q^0.

Brown and Weikard [BW] considered the Schrödinger operator $-y'' + (p_A + q)y$ on the half-line, where p_A is an algebro-geometric potentials and q is a compactly supported potential. They proved that the zeros of the Jost function determine q uniquely.

Christiansen [Ch] considered resonances associated to the Schrödinger operator $-y'' + (p_S + q)y$ on the real line, where p_S is a step potential. She determined asymptotics of the resonance-counting function. Moreover, she obtained that the resonances determine q uniquely.



FIGURE 2. The cut domain $\mathcal{Z} = \mathbb{C} \setminus \cup g_n$ and the slits $g_n = (e_n^-, e_n^-)$ in the z-plane.

In the proof of theorems we use properties of the quasimomentum from [KK], [M], a priori estimates from [KK], [M], and results from the inverse theory for the Hill operator from [K5], see (4.2).

The plan of the paper is as follows. In Section 2 we describe the preliminary results about fundamental solutions. In Sections 3 we study the function ξ and prove Theorem 1.1. In Sections 4 we prove the main Theorem 1.2-1.3, which are crucial for Section 3 and 4.

2. PRELIMINARIES

We will work with the momentum $z = \sqrt{\lambda}$, where λ is an energy and recall that $\sqrt{1} = 1$. If $\lambda \in \gamma_n, n \geqslant 1$, then $z \in g_{\pm n}$ and if $\lambda \in \gamma_0 = (-\infty, 0)$, then $z \in g_0^\pm = i\mathbb{R}_\pm$, where the momentum gaps g_n are given by

$$g_n = (e_n^-, e_n^+) = -g_{-n}, \quad e_n^\pm = \sqrt{E_n^\pm} > 0, \quad n \geqslant 1, \quad \textit{and} \quad \Delta(e_n^\pm) = (-1)^n. \tag{2.1}$$

Introduce the cut domain (see Fig.2)

$$\mathcal{Z}_0 = \mathbb{C} \setminus \cup \overline{g}_n.$$

Below we will use the momentum variable $z = \sqrt{\lambda}$ and the corresponding the Riemann surface $\mathcal{M}$, which is more convenient for us, than the Riemann surface Λ. Slitting the n-th momentum gap g_n (suppose it is nontrivial) we obtain a cut G_n with an upper g_n^+ and lower rim g_n^-. Below we will identify this cut G_n and the union of of the upper rim (gap) $\overline{g}_n^+$ and the lower rim (gap) $\overline{g_n}^-$, i.e.,

$$G_n = \overline{g}_n^+ \cup \overline{g}_n^-, \quad \textit{where} \quad g_n^\pm = g_n \pm i0. \tag{2.2}$$

Also we will write $z \pm i0 \in g_n^\pm$, if $z \in g_n$.

In order to construct the Riemann surface $\mathcal{M}$ we take the cut domain $\mathcal{Z}_0 = \mathbb{C} \setminus \cup \overline{g}_n$ and identify (i.e. we glue) the upper rim g_n^+ of the slit G_n with the upper rim g_{-n}^+ of the slit G_{-n} and correspondingly the lower rim g_n^- of the slit G_n with the lower rim g_{-n}^- of the slit G_{-n} for all nontrivial gaps.

The mapping $z = \sqrt{\lambda}$ from Λ onto $\mathcal{M}$ is one-to-one and onto. The gap $\gamma_n^{(1)} \subset \Lambda_1$ maps onto $g_n^+ \subset \mathcal{M}_1$ and the gap $\gamma_n^{(2)} \subset \Lambda_2$ maps onto $g_n^- \subset \mathcal{M}_2$. From a physical point of view, the upper rim g_n^+ is a physical gap and the lower rim g_n^- is a non physical gap. Moreover, $\mathcal{M} \cap \mathbb{C}_+ = \mathcal{Z}_0 \cap \mathbb{C}_+$ plus all physical gaps g_n^+ is a so-called physical "sheet" $\mathcal{M}_1$ and $\mathcal{M} \cap \mathbb{C}_- = \mathcal{Z}_0 \cap \mathbb{C}_-$ plus all non physical gaps g_n^- is a so-called non physical "sheet" $\mathcal{M}_2$. The set (the spectrum) $\mathbb{R} \setminus \cup g_n$ joints the first and second sheets.



FIGURE 3. The domain $\mathcal{K} = \mathbb{C} \setminus \cup\Gamma_n$, where the slit $\Gamma_n = (\pi n - ih_n, \pi n + ih_n)$

We introduce the quasimomentum $k(\cdot)$ for H_0 by $k(z) = \arccos \Delta(z),\ z \in \mathcal{Z}_0 = \mathbb{C} \setminus \cup \overline{g}_n$. The function $k(z)$ is analytic in $z \in \mathcal{Z}_0$ and satisfies:

$$(i) \quad k(z) = z + O(1/z) \quad as \quad |z| \to \infty, \quad and \quad (ii) \quad \operatorname{Re} k(z \pm i0)|_{[e_n^-, e_n^+]} = \pi n, \quad n \in \mathbb{Z}, \tag{2.3}$$

and $\pm \operatorname{Im} k(z) > 0$ for any $z \in \mathbb{C}_\pm$, see [M], [KK]. The function $k(\cdot)$ is analytic on $\mathcal{M}$ and satisfies $\sin k(z) = \Omega(z^2), z \in \mathcal{M}$. Moreover, the quasimomentum $k(\cdot)$ is a conformal mapping from $\mathcal{Z}_0$ onto the quasimomentum domain $\mathcal{K} = \mathbb{C} \setminus \cup\Gamma_n$, where the slit $\Gamma_n = (\pi n - ih_n, \pi n + ih_n)$. The height h_n is defined by the equation $\cosh h_n = (-1)^n \Delta(e_n)$, where $e_n \in [e_n^-, e_n^+]$ and $\Delta'(e_n) = 0$. The Floquet solutions $\psi_\pm(x, z), z \in \mathcal{Z}_0$ of H_0 is given by

$$\psi_\pm(x, z) = \vartheta(x, z) + m_\pm(z)\varphi(x, z), \quad m_\pm = \frac{\beta \pm i \sin k}{\varphi(1, \cdot)}, \quad \beta = \frac{\varphi'(1, \cdot) - \vartheta(1, \cdot)}{2}, \tag{2.4}$$

where $\varphi(1, z)\psi^+(\cdot, z) \in L^2(\mathbb{R}_+)$ for all $z \in \mathbb{C}_+ \cup \cup g_n$. If $p = 0$, then $k = z$ and $\psi_\pm(x, z) = e^{\pm izx}$.

The function $\sin k$ and each function $\varphi(1, \cdot)\psi_\pm(x, \cdot), x \in \mathbb{R}$ are analytic on the Riemann surface $\mathcal{M}$. Recall that the Floquet solutions $\psi_\pm(x, z), (x, z) \in \mathbb{R} \times \mathcal{M}$ satisfy (see [T])

$$\psi_\pm(0, z) = 1, \quad \psi_\pm(0, z)' = m_\pm(z), \quad \psi_\pm(1, z) = e^{\pm ik(z)}, \quad \psi_\pm(1, z)' = e^{\pm ik(z)} m_\pm(z), \tag{2.5}$$

$$\psi_\pm(x, z) = e^{\pm ik(z)x}(1 + O(1/z)) \quad as \quad |z| \to \infty, \quad (x, z) \in \mathbb{R} \times \mathcal{Z}_\varepsilon, \tag{2.6}$$

where the set $\mathcal{Z}_\varepsilon = \{z \in \mathcal{Z}_0 : \operatorname{dist}\{z, g_n\} > \varepsilon, g_n \neq \emptyset, n \in \mathbb{Z}\},\ \varepsilon > 0$. Below we need the simple identities

$$\beta^2 + 1 - \Delta^2 = 1 - \varphi'(1, \cdot)\vartheta(1, \cdot) = -\varphi(1, \cdot)\vartheta'(1, \cdot). \tag{2.7}$$

Let $\mathcal{D}_r(z_0) = \{|z - z_0| < r\}$ be a disk for some $r > 0$. Recall the well known properties of the function $m_\pm$ (see [T]).

Lemma 2.1. *i)* $\operatorname{Im} m_+(z) > 0$ *for all* $(z, n) \in (z_{n-1}^+, z_n^-) \times \mathbb{N}$ *and the following asymptotics hold true:*

$$m_\pm(z) = \pm iz + O(1) \quad as \quad |z| \to \infty, \quad z \in \mathcal{Z}_\varepsilon, \varepsilon > 0. \tag{2.8}$$

ii) Let $g_n = \emptyset$ *for some* $n \in \mathbb{Z}$. *Then the functions* $\sin k(\cdot), m_\pm$ *are analytic in the disk* $B(\mu_n, \varepsilon) \subset \mathcal{Z}_0$ *for some* $\varepsilon > 0$ *and the functions* $\sin k(z)$ *and* $\varphi(1, z)$ *have the simple zero at* μ_n. *Moreover,* $m_\pm$ *satisfies*

$$m_\pm(\mu_n) = \frac{\beta'(\mu_n) \pm i(-1)^n k'(\mu_n)}{\partial_z \varphi(1, \mu_n)}, \quad \operatorname{Im} m_\pm(\mu_n) \neq 0. \tag{2.9}$$

iii) If the function m_+ *has a pole at* $\mu_n + i0$ *for some* $n \in \mathbb{N}$*, then* $k(\mu_n + i0) = \pi n + ih_{sn}$ *and*

$$h_{sn} > 0, \quad \beta(\mu_n) = i \sin k(\mu_n + i0) = -(-1)^n \sinh h_{sn}, \quad m_+ \in \mathscr{A}(\mu_n - i0),$$

$$m_+(\mu_n + z) = \frac{\rho_n + O(z)}{z} \quad \text{as } z \to 0, \ z \in \mathbb{C}_+, \qquad \rho_n = \frac{-2\sinh|h_{sn}|}{(-1)^n \partial_z \varphi(1, \mu_n)} < 0. \tag{2.10}$$

iv) If the function m_+ *has a pole at* $\mu_n - i0$ *for some* $n \in \mathbb{N}$*, then* $k(\mu_n - i0) = \pi n + ih_{sn}$ *and*

$$h_{sn} < 0, \quad \beta(\mu_n) = -i \sin k(\mu_n - i0) = (-1)^n \sinh h_{sn}, \ m_+ \in \mathscr{A}(\mu_n^+),$$

$$m_+(\mu_n + z) = \frac{\rho_n + O(z)}{z} \quad \text{as } z \to 0, \ z \in \mathbb{C}_-. \tag{2.11}$$

iv) $\mu_n = e_n^-$ *or* $\mu_n = e_n^+$ *(here* $e_n^- \neq e_n^+$*) for some* $n \neq 0$ *iff*

$$m_+(\mu_n + z) = \frac{\rho_n^\pm + O(z)}{\sqrt{z}} \quad \text{as } z \to 0, \ z \in \mathbb{C}_+, \qquad \text{some const } \rho_n^\pm \neq 0. \tag{2.12}$$

The following asymptotics hold true as $n \to \infty$ (see [PT], [K5]):

$$\mu_n = \pi n + \varepsilon_n(p_{c0} - p_{cn} + O(\varepsilon_n)), \qquad \varepsilon_n = 1/2\pi n, \tag{2.13}$$

$$e_n^\pm = \pi n + \varepsilon_n(p_0 \pm |p_n| + O(\varepsilon_n)), \qquad p_n = \int_0^1 p(x) e^{-i2\pi n x} dx = p_{cn} - ip_{sn}. \tag{2.14}$$

The equation $-f'' + (p+q)f = z^2 f, \ z \neq e_n^\pm$ has unique solutions $f_\pm(x,z)$ such that

$$f_+(x,z) = \psi_+(x,z), \ x \geqslant t, \qquad \text{and} \qquad f_-(x,z) = \psi_-(x,z), \ x \leqslant 0, \tag{2.15}$$

and $f_+(x,z) = \overline{f}_+(x,-z), z^2 \in \sigma(H_0) \setminus \{e_n^\pm, n \in \mathbb{Z}\}$. This yields

$$f_+(x,z) = b(z) f_-(x,z) + a(z) f_-(x,-z), \quad a = \frac{w}{w_0}, \quad b = \frac{s}{w_0}, \tag{2.16}$$

where $z^2 \in \sigma(H_0) \setminus \{e_n^\pm, n \in \mathbb{Z}\}$ and

$$w = \{f_-, f_+\}, \qquad s = \{f_+(x,z), f_-(x,-z)\}, \quad w_0 = \{\psi_-, \psi_+\} = \frac{2i \sin k}{\varphi(1, \cdot)} \tag{2.17}$$

and $\{f, g\} = fg' - f'g$ is the Wronskian. The scattering matrix $\mathcal{S}_M$ for H, H_0 is given by

$$\mathcal{S}_M(z) \equiv \begin{pmatrix} a(z)^{-1} & r_-(z) \\ r_+(z) & a(z)^{-1} \end{pmatrix}, \qquad r_\pm = \frac{s(\mp z)}{w(z)} = \mp \frac{b(\mp z)}{a(z)}, \qquad z^2 \in \sigma_{ac}(H), \tag{2.18}$$

where $1/a$ is the transmission coefficient and $r_\pm$ is the reflection coefficient. We have the following identities from [F1], [F3]:

$$|a(z)|^2 = 1 + |b(z)|^2, \qquad z^2 \in \sigma_{ac}(H), \tag{2.19}$$

$$a(z) = D(z^2), \qquad z \in \mathcal{Z}_0. \tag{2.20}$$

The functions a, b, s, w are analytic in $\mathcal{Z}_0$ and real on $i\mathbb{R}$. Then the following identities hold true:

$$a(-z) = \overline{a}(\overline{z}), \quad w(-z) = \overline{w}(\overline{z}), \quad s(-z) = \overline{s}(\overline{z}), \quad w_0(-z) = \overline{w}_0(\overline{z}), \quad z \in \mathcal{Z}_0. \tag{2.21}$$

Let $\varphi(x, z, \tau), \ (z, \tau) \in \mathbb{C} \times \mathbb{R}$ be the solutions of the equation

$$-\varphi'' + p(x+\tau)\varphi = z^2 \varphi, \qquad \varphi(0, z, \tau) = 0, \quad \varphi'(0, z, \tau) = 1. \tag{2.22}$$

The function $\varphi(1, z, x)$ for all $(x, z) \in R \times \mathbb{C}$ satisfies the following identity (see [Tr])

$$\varphi(1, \cdot, \cdot) = \varphi(1, \cdot)\vartheta^2 - \vartheta'(1, \cdot)\varphi^2 + 2\beta\varphi\vartheta = \varphi(1, \cdot)\psi_-\psi_+. \tag{2.23}$$

Let $\widetilde{\vartheta}, \widetilde{\varphi}$ be the solutions of the equations $-y'' + (p + q)y = z^2 y, z \in \mathbb{C}$ and satisfying

$$\widetilde{\varphi}(x, z) = \varphi(x, z), \qquad \widetilde{\vartheta}(x, z) = \vartheta(x, z) \quad for \ \ all \ x \geqslant t. \tag{2.24}$$

A solution of the equation $-y'' + (p - z^2)y = f, y(0) = y'(0) = 0$ has the form $y = \int_0^x \varphi(x - \tau, z, \tau)f(\tau)d\tau$. Hence the solutions $\widetilde{\vartheta}, \widetilde{\varphi}$ and f_+ of the equation $-y'' + (p + q)y = z^2 y$ satisfy the equation

$$y(x, z) = y_0(x, z) - \int_x^t \varphi(x - \tau, z, \tau)q(\tau)y(\tau, z)d\tau, \qquad x \leqslant t, \tag{2.25}$$

where y is one from $\widetilde{\vartheta}, \widetilde{\varphi}$ and f_+; y_0 is the corresponding function from ϑ, φ and ψ_+. For each $x \in \mathbb{R}$ the functions $\widetilde{\vartheta}(x, z), \widetilde{\varphi}(x, z)$ and $\vartheta(x, z), \varphi(x, z)$ are entire in $z \in \mathbb{C}$ and satisfy

$$\max\{||z|_1\widetilde{\varphi}(x, z)|, \ |\widetilde{\varphi}'(x, z)|, |\widetilde{\vartheta}(x, z)|, \frac{1}{|z|_1}|\widetilde{\vartheta}'(x, z)|\} \leqslant X_1 = e^{|\operatorname{Im} z||2t-x|+\|q\|_t+\|p\|_t+\int_x^t |p(\tau)|d\tau},$$

$$|z|_1 = \max\{1, |z|\}, \qquad |\widetilde{\vartheta}(x, z) - \vartheta(x, z)| \leqslant \frac{X_1}{|z|}\|q\|_t, \quad |\widetilde{\varphi}(x, z) - \varphi(x, z)| \leqslant \frac{X_1}{|z|^2}\|q\|_t, \tag{2.26}$$

where $\|p\|_t = \int_0^t |p(s)|ds$ and

$$\max\{|z|_1|\varphi(x, z)|, \ |\varphi'(x, z)|, |\vartheta(x, z)|, \frac{1}{|z|_1}|\vartheta'(x, z)|\} \leqslant X = e^{|\operatorname{Im} z|x+\|p\|_x},$$

$$|\varphi(x, z) - \frac{\sin zx}{z}| \leqslant \frac{X}{|z|^2}\|p\|_x, \quad |\vartheta(x, z) - \cos zx| \leqslant \frac{X}{|z|}\|p\|_x, \tag{2.27}$$

for $(p, x, z) \in L^1_{loc}(\mathbb{R}) \times \mathbb{R} \times \mathbb{C}$, see [PT]. These estimates yield

$$\beta(z) = \int_0^1 \frac{\sin z(2x - 1)}{z}p(x)dx + \frac{O(e^{|\operatorname{Im} z|})}{z^2},$$

$$\beta'(z) = \int_0^1 \frac{\cos z(2x - 1)}{z}p(x)(2x - 1)dx + \frac{O(e^{|\operatorname{Im} z|})}{z^2} \quad as \quad |z| \to \infty. \tag{2.28}$$

Lemma 2.2. *For all $z \in \mathcal{Z}_0$ the following identities and asymptotics hold true:*

$$f_+(\cdot, z) = \widetilde{\vartheta}(\cdot, z) + m_+(z)\widetilde{\varphi}(\cdot, z), \tag{2.29}$$

$$f_+(0, z)= 1+\int_0^t \varphi(x, z)q(x)f_+(x, z)dx, \ \ f_+'(0, z)= m_+(z) - \int_0^t \vartheta(x, z)q(x)f_+(x, z)dx, \tag{2.30}$$

$$f_+(x, z) = e^{ik(z)x} + e^{(2t-x)\mathfrak{J}}O(1/z), \qquad \mathfrak{J} = \frac{|\operatorname{Im} z| - \operatorname{Im} z}{2}, \quad x \in [0, t], \tag{2.31}$$

$$f_+(0, z) = 1 + e^{2t\mathfrak{J}}O(1/z), \quad f_+'(0, z) = iz + O(1) + e^{2t\mathfrak{J}}o(1) \tag{2.32}$$

as $|z| \to \infty, z \in \mathcal{Z}_\varepsilon, \varepsilon > 0$, where $\widehat{q}(z) = \int_0^t q(x)e^{2izx}dx, \ z \in \mathbb{C}$.

Proof. Using (2.24), (2.15) we obtain (2.29). Using the identity $\varphi(x,\cdot,s)=\vartheta(t,\cdot)\varphi(x+s,\cdot)-\varphi(s,,\cdot)\vartheta(x+s,\cdot)$, we obtain $\varphi(-s,\cdot,s)=-\varphi(s,\cdot)$ and $\varphi'(-s,\cdot,s)=\vartheta(s,\cdot)$. Substituting the last identities into (2.25) we get (2.30).

Standard iteration arguments (see [PT]) for the equation (2.25) give (2.31).

Substituting (2.31) into (2.25) we obtain (2.32). ■

Lemma 2.3. *i) For each* $z\in\mathcal{Z}_0$ *the following identities hold true:*

$$w(z)=f_+'(0,z)-m_-(z)f_+(0,z)=w_0(z)-\int_0^t q(x)\psi_-(x,z)f_+(x,z)dx, \tag{2.33}$$

$$s(z)=f_+(0,z)m_+(z)-f_+'(0,z)=\int_0^t q(x)\psi_+(x,z)f_+(x,z)dx. \tag{2.34}$$

The functions $\xi(z)=2i\sin k(z)a(z)$ *and* $s(\cdot)$ *have the following asymptotics:*

$$\xi(z)=2i\sin z(1+O(e^{2t\Im}/z)),\qquad s(z)=O(e^{2t\Im}), \tag{2.35}$$

as $|z|\to\infty, z\in\mathcal{Z}_\varepsilon,\varepsilon>0$.

ii) The function $s(\cdot)$ *has exponential type* $\rho_\pm$ *in the half plane* $\mathbb{C}_\pm$*, where* $\rho_+=0,\rho_-=2t$.

Proof. We have $w=\{f^-,f^+\}=\psi^-f^{+\prime}-m^-f^+|_{x=0}$, which yields the identity in (2.33). Substituting (2.30) into (2.33) we obtain (2.34). Asymptotics from Lemma 2.2 and (2.8) imply (2.35).

ii) We show $\rho_-=2t$. Due to (2.35), s has exponential type $\rho_-\leqslant 2t$. The decompositions $f_+=e^{ixz}(1+h)$ and $\psi_+=e^{ixz}(1+\eta)$ give $(1+h)(1+\eta)=1+T, T=h+\eta+\eta h$ and

$$s(z)=\int_0^t q(x)\psi^+(x,z)f^+(x,z)dx=\int_0^t q(x)e^{i2xz}(1+T(x,z))dx,\qquad z\in\mathcal{Z}_\varepsilon. \tag{2.36}$$

Asymptotics (2.6), (2.27), (2.31) and $k(z)=z+O(1/z)$ as $|z|\to\infty$ (see [KK]) yield

$$\eta(x,z)=O(1/z),\qquad h(x,z)=e^{2(t-x)|\operatorname{Im} z|}O(1/z)\quad as\ |z|\to\infty,\quad z\in\mathcal{Z}_\varepsilon. \tag{2.37}$$

We need the following variant of the Paley Wiener type Theorem from [Fr]: *let* $q\in\mathcal{Q}_t$ *and let each* $G(x,z),x\in[0,t]$ *be analytic for* $z\in\mathbb{C}_-$ *and* $G\in L^2((0,t)dx,\mathbb{R}dz)$. *Then* $\int_0^t e^{2izx}q(x)(1+G(x,z))dx$ *has exponential type at least* $2t$ *in* $\mathbb{C}_-$.

We can not apply this result to the function $T(x,z), z\in\mathbb{C}_-$, since $m_+(z)$ may have a singularity at $\mu_n-i0\in\overline{g}_n^-$ if $g_n\neq\emptyset$. But we can use this result for the function $T(x,z-i), z\in\mathbb{C}_-$, since (2.36), (2.37) imply $\sup_{x\in[0,1]}|T(x,-i+\tau)|=O(1/\tau)$ as $\tau\to\pm\infty$. Then the function $s(z)$ has exponential type $2t$ in the half plane $\mathbb{C}_-$. The proof for $\rho_+=0$ is similar. ■

3. ANALYZE OF THE FUNCTION ξ

Below we need the identities and the asymptotics as $n\to\infty$ from [KK]:

$$(-1)^{n+1}i\sin k(z)=\sinh v(z)=\pm|\Delta^2(z)-1|^{\frac12}>0\quad all\quad z\in g_n^\pm, \tag{3.1}$$

$$v(z)=\pm|(z-e_n^-)(e_n^+-z)|^{\frac12}(1+O(n^{-2})),\quad \sinh v(z)=v(z)(1+O(|g_n|^2)),\quad z\in\overline{g}_n^\pm. \tag{3.2}$$

Let $\nu_n^2, n\geqslant 1$ be the Neumann spectrum of the equation $-y''+py=\nu^2y$ on the interval $[0,1]$ with the boundary condition $y(0)=y(1)=0$. It is well known that each $\nu_n^2\in[E_n^-,E_n^+], n\geqslant 1$.

Lemma 3.1. *Let $p \in L^1(0,1)$.*
i) Then the following asymptotics hold true uniformly for $z \in [e_n^-, e_n^+]$ as $n \to \infty$:

$$\varphi(1,z) = (-1)^n \frac{(z-\mu_n)}{\pi n}(1+O(1/n)), \tag{3.3}$$

$$-\frac{\vartheta'(1,z)}{z^2} = (-1)^n \frac{(z-\nu_n)}{\pi n}(1+O(1/n)). \tag{3.4}$$

ii) Let $z \in g_n$ and $e_n^- \geqslant R_p = 8e^{\|p\|_1}$. Then the following estimates hold true (here $\dot{u} = \partial_z u$)

$$|g_n|^2 \leqslant \frac{8e^{\|p\|_1}}{|z_n|} < 1, \tag{3.5}$$

$$|\dot{\varphi}(1,z)| \leqslant \frac{3e^{\|p\|_1}}{2|z|}, \qquad |\varphi(1,z)|_{z\in g_n} \leqslant |g_n|\frac{3e^{\|p\|_1}}{2|z|}, \tag{3.6}$$

$$|\dot{\vartheta}'(1,z)| \leqslant |z|\frac{3e^{\|p\|_1}}{2}, \qquad |\vartheta'(1,z)| \leqslant |g_n||z|\frac{3e^{\|p\|_1}}{2}, \tag{3.7}$$

$$|\dot{\beta}(z)| \leqslant \frac{3e^{\|p\|_1}}{2|z|}, \qquad |\beta(e_n^\pm)| \leqslant |g_n|\frac{3e^{\|p\|_1}}{2}, \qquad |\beta(z)| \leqslant |g_n|\frac{9e^{\|p\|_1}}{4|z|}. \tag{3.8}$$

iii) In each disk $\mathcal{D}_{\frac{\pi}{4}}(\pi n) \subset \mathcal{D} = \{|z| > 32e^{2\|p\|_1}\}$ there exists exactly one momentum gap g_n of the operator H_0. Moreover, if $g_n, g_{n+1} \subset \mathcal{D}$, then $e_{n+1}^- - e_n^+ \geqslant \pi$.

Proof. i) We have the Taylor series $\varphi(1,z) = \dot{\varphi}(1,\mu_n)\tau + \ddot{\varphi}(1,\mu_n+\alpha\tau)\frac{\tau^2}{2}$ for any $z \in [e_n^-, e_n^+]$ and some $\alpha \in [0,1]$, where $\tau = z - \mu_n$ and $\dot{\varphi} = \partial_z\varphi$. Asymptotics (2.27) give $\dot{\varphi}(1,\mu_n) = 2(-1)^n(1+O(1/n))/(2\pi n)$ and $\ddot{\varphi}(1,\mu_n+\alpha\tau)\tau = O(n^{-2})$, which yields (3.3). Similar arguments imply (3.4).

ii) Using $|\Delta(z) - \cos z| \leqslant \frac{e^{\|p\|_1}}{|z|}$, for all $|z| \geqslant 2$, we obtain

$$\frac{h_n^2}{2} \leqslant \cosh h_n - 1 = |\Delta(z_n)| - 1 \leqslant \frac{e^{\|p\|_1}}{|z_n|}.$$

Then the estimate $|g_n| \leqslant 2h_n$ (see [KK]) gives (3.5).

Due to (2.27), the function $f = z\varphi(1,z)$ has the estimate $|f(z)| \leqslant C_0 = e^{\|p\|_1}, z \in \mathbb{R}$. Then the Bernstein inequality gives $|\dot{f}(z)| = |\varphi(1,z) + z\dot{\varphi}(1,z)| \leqslant C_0, z \in \mathbb{R}$, which yields $|\dot{\varphi}(1,z)| \leqslant \frac{C_0}{|z|}(1+\frac{1}{|z|}), z \in \mathbb{R}$. Moreover, we obtain $|\varphi(1,z)| \leqslant |g_n| \max_{z\in g_n} |\dot{\varphi}(1,z)| \leqslant |g_n|\frac{3C_0}{2|z|}$.

The proof of (3.7) and the estimate $|\dot{\beta}(z)| \leqslant \frac{3e^{\|p\|_1}}{2|z|}$ is similar. Identity (2.7) gives $\beta^2(e_n^\pm) = -\varphi(1,e_n^\pm)\vartheta'(1,e_n^\pm)$. Then (3.6), (3.7) imply $|\beta(e_n^\pm)| \leqslant |g_n|\frac{3e^{\|p\|_1}}{2}$. Using these estimates and $\beta(z) = \beta(e_n^-) + \beta'(z_*)(z - e_n^-)$ for some $z_* \in g_n$ we obtain (3.8).

iii) Using (2.27) we obtain

$$|(\Delta^2(z) - 1) - (\cos^2 z - 1)| \leqslant 2X|\Delta(z) - \cos z| \leqslant 2X^2/|z|, \quad X = e^{|\operatorname{Im}|+\|p\|_1}.$$

After this the standard arguments (due to Rouche's theorem) give the proof of iii). ■

The function a is not convenient, since a is not analytic on $\mathcal{M}$. The modified function ξ in the momentum variable z is given by

$$\xi(z) = 2i\sin k(z)a(z) = \Xi(z^2), \qquad z \in \mathcal{Z}_0. \tag{3.9}$$

The number $e_n = \sqrt{E_n} \in [e_n^-, e_n^+]$ satisfies $|\Delta(e_n)| = \max_{z\in g_n} |\Delta(z)| = \cosh h_n \geqslant 1$ for some $h_n \geqslant 0$.

Lemma 3.2. *i) The following identity and asymptotics hold true:*

$$\xi(z)=2i\sin k(z)(1-A(z^2))-J(z^2),\quad z\in\mathcal{M},\qquad J(z^2)=\int_{\mathbb{R}}q(x)Y_1(x,z)dx$$

$$A(z^2)=\int_{\mathbb{R}}q(x)Y_2(x,z)dx,\qquad Y_1=\varphi_1\vartheta\widetilde{\vartheta}-\vartheta_1'\varphi\widetilde{\varphi}+\beta(\varphi\widetilde{\vartheta}+\vartheta\widetilde{\varphi}),\qquad Y_2=\frac{1}{2}(\varphi\widetilde{\vartheta}-\vartheta\widetilde{\varphi}),$$

$$\xi(z)=2(-1)^{n+1}(1+A(z^2))\sinh v(z)-J(z^2),\qquad z\in g_n^{\pm}\neq\emptyset,\quad (3.10)$$

where $v=\operatorname{Im}k$ *and* $\pm v(z)>0$ *for* $z\in g_n^{\pm}$*,*

$$Y_1=\varphi(1,z,x)+Y_{11}(z,x),\quad Y_{11}=\varphi_1\vartheta\widetilde{\vartheta}_*-\vartheta_1'\varphi\widetilde{\varphi}_*+\beta(\varphi\widetilde{\vartheta}_*+\vartheta\widetilde{\varphi}_*),\quad \widetilde{\vartheta}_*=\widetilde{\vartheta}-\vartheta,\widetilde{\varphi}_*=\widetilde{\varphi}-\varphi,$$

$$Y_{11}(z,x)=O(|g_n|/n^2)\quad as\quad z\in g_n,n\to\infty.\quad (3.11)$$

Moreover, the functions J,A *are entire and* ξ *is analytic on* $\mathcal{M}$*.*
ii) Let $|z|\geqslant 2$*. Then the following estimates hold true*

$$|J(z^2)|\leqslant C_{p,q}\|q\|_t\biggl(|\varphi(1,z)|+\frac{|\vartheta'(1,z)|}{|z|^2}+\frac{|\beta(z)|}{|z|}\biggr)e^{2t|\operatorname{Im}z|}\leqslant\frac{3C_*}{|z|}e^{(2t+1)|\operatorname{Im}z|},$$

$$|A(z^2)|\leqslant\frac{\|q\|_t^2C_{p,q}}{|z|^2}e^{2t|\operatorname{Im}z|},\quad where\quad C_*=\|q\|_te^{\|p\|_1+\|q\|_t+2\|p\|_t},\quad C_{p,q}=e^{2\|p\|_t+\|q\|_t},\quad (3.12)$$

$$|J(e_n^2)|\leqslant\frac{R_1}{|z_n|}\sinh h_n,\qquad if\quad e_n^-\geqslant R_p=8e^{\|p\|_1},\qquad R_1=11C_*,\qquad (3.13)$$

Proof. i) We rewrite the identity (2.34) in the form

$$\xi(z)=2i\sin k(z)-\int_{\mathbb{R}}q(x)Y(x,z)dx,\quad Y=\varphi_1\psi_-(x,z)f_+(x,z),\qquad (3.14)$$

for $z\in\mathbb{C}_+$. Using (3.14), (2.29) we rewrite Y in the form

$$Y=\varphi_1(\vartheta+m_-\varphi)(\widetilde{\vartheta}+m_+\widetilde{\varphi})=\varphi_1\biggl(\vartheta\widetilde{\vartheta}+m_+\vartheta\widetilde{\varphi}+m_-\varphi\widetilde{\vartheta}-\frac{\vartheta_1'}{\varphi_1}\varphi\widetilde{\varphi}\biggr)=Y_1-i2Y_2\sin k.$$

Substituting the last identity into $\int_{\mathbb{R}}q(x)Y(x,z)dx$ and using (3.1), we obtain (3.10).

Substituting asymptotics from (2.26), Lemma 3.1 into Y_{11} we obtain $Y_{11}(z,x)=O(|g_n|/n^2)$ as $z\in g_n,n\to\infty$.

ii) Using (2.26), (2.27) and (3.10) and Lemma 3.1, we obtain (3.12). Estimates (3.12) and Lemma 3.1 give $|J(e_n^2)|\leqslant\frac{21C_*}{4|z_n|}|g_n|$; and the estimate $|g_n|\leqslant 2|h_n|\leqslant 2\sinh h_n$ from [KK] yields (3.13). ■

This lemma gives that the function ξ is analytic on $\mathcal{M}$ and the function Ξ is analytic on Λ. We define the bound states, resonances in terms of momentum variable $z\in\mathcal{M}$. Recall that there are bound states on the physical gaps and resonances on the non physical gaps. Define the set $\mathcal{M}_0=\{z\in\mathcal{M}:z=e_{\pm n}^+\in\mathcal{M}_1$ and $z=e_{\pm n}^+\in\mathcal{M}_2,\gamma_n=\emptyset,n\geqslant 1\}\subset\mathcal{M}$. The set $\mathcal{M}_0$ is the image of Λ_0 (see before Definition of states, Section 1) under the mapping $z=\sqrt{\lambda}$.

Definition. *Each zero of* $\xi(z)=2i\sin k(z)a(z)$ *in* $\mathcal{M}\setminus\mathcal{M}_0$ *is a state of* H*.*
1) A state $z\in\mathcal{M}_2$ *is a resonance.*
2) A state $z=e_n^{\pm},n\in\mathbb{Z}$ *is a virtual state.*
3) A resonance $z\in\cup g_n^-\subset\mathcal{M}_2$ *is an anti-bound state.*

Of course, z^2 is really the energy, but since the momentum z is the natural parameter, we will abuse the terminology.

The kernel of the resolvent $R = (H - z^2)^{-1}, z \in \mathbb{C}_+$, has the form

$$R(x, x', z) = \frac{f_-(x, z) f_+(x', z)}{-w(z)} = \frac{R_1(x, x', z)}{-\xi(z)}, \quad x < x', \quad R_1 = \varphi(1, z) f_-(x, z) f_+(x', z),$$

and $R(x, x', z) = R(x', x, z),\ x > x'$. Identity (2.29) yields $f_\pm = \widetilde{\vartheta} + m_\pm \widetilde{\varphi}, \quad \widetilde{\vartheta} = \widetilde{\vartheta}(x), \quad \widetilde{\varphi} = \widetilde{\varphi}(x)$. Let $\widetilde{\vartheta}_* = \widetilde{\vartheta}(x'), \quad \widetilde{\varphi}_* = \widetilde{\varphi}(x')$. Then using (2.7) we obtain

$$R_1(x, x', z) = \varphi(1, \cdot) \widetilde{\vartheta} \widetilde{\vartheta}_* + (\beta - i \sin k) \widetilde{\varphi} \widetilde{\vartheta}_* + (\beta + i \sin k) \widetilde{\vartheta} \widetilde{\varphi}_* - \vartheta'(1, \cdot) \widetilde{\varphi}_* \widetilde{\varphi}.$$

Thus if $\xi(z) = \varphi(1, z) w(z) = 0$ at some $z \in \mathcal{M}$, then $(H - z^2)^{-1}$ has singularity at z. The poles of $R(x, x', z)$ define the bound states and resonances. The zeros of ξ define the bound states and resonances, since the function $R_1 = \varphi_1(z) f^-(x, z) f^+(x', z)$ is locally bounded.

If $q = 0$, then $R_0 = (H_0 - z^2)^{-1}$ has the form

$$R_0(x, x', z) = \frac{R_{10}(x, x', z)}{-\xi_0(z)}, \quad \xi_0 = \varphi(1, z) w_0(z) = 2i \sin k(z),$$

$$R_{10} = \varphi_1(z) \psi_-(x, z) \psi_+(x', z) = \varphi(1, \cdot) \vartheta \vartheta_* + (\beta - i \sin k) \varphi \vartheta_* + (\beta + i \sin k) \vartheta \varphi_* - \vartheta'(1, \cdot) \varphi \varphi_*,$$

where $\varphi = \varphi(x, z), ..$ and $\varphi_* = \varphi(x', z), ..$ Thus $R_0(x, x', z)$ has singularity at some $z \in \mathcal{M}$ iff $\sin k(z) = 0$, i.e., $k(z) = \pi n$ and then $z = e_n^\pm$.

Define the functions F, S by

$$F = \xi(z) \xi(-z), \qquad S(z) = \varphi^2(1, z) s(z) s(-z), \quad z \in \mathcal{Z}_0.$$

Lemma 3.3. *i) The functions $F(z), S(z), z \in \mathcal{Z}_0$ have analytic continuations into the whole complex plane $\mathbb{C}$ and satisfy*

$$F(z) = 4(1 - \Delta^2(z))(1 + A(z^2))^2 + J^2(z^2) = 4(1 - \Delta^2(z^2)) + S(z), \qquad z \in \mathbb{C}. \tag{3.15}$$

Moreover, $F(z) > 0$ and $S(z) \geqslant 0$ on each interval $(e_{n-1}^+, e_n^-), n \geqslant 1$ and F has even number of zeros on each interval $[e_n^-, e_n^+], n \geqslant 1$. The function F has only simple zeros at $e_n^\pm, g_n \neq \emptyset$.
ii) If $g_n = (e_n^-, e_n^+) = \emptyset$ for some $n \neq 0$, then each $f_\pm(x, \cdot), x \in \mathbb{R}$ is analytic in some disk $\mathcal{D}(e_n^+, \varepsilon), \varepsilon > 0$. Moreover, $\mu_n = e_n^\pm$ is a double zero of F and e_n^+ is not a state of H.
iii) Let $z \in g_n^+$ be a bound state for some $n \geqslant 1$, i.e., $\xi(z + i0) = 0$. Then $z - i0 \in g_n^-$ is not an anti-bound state and $\xi(z - i0) \neq 0$.
iv) Let $z \in i\mathbb{R}_+$ be a bound state, i.e., $\xi(z) = 0$. Then $-z \in i\mathbb{R}_-$ is not an anti-bound state and $\xi(-z) \neq 0$.
v) $z \in \mathbb{C}_- \setminus i\mathbb{R}$ is a zero of F iff $z \in \mathbb{C}_- \setminus i\mathbb{R}$ is a zero of ξ (with the same multiplicity).
vi) $z \in i\mathbb{R}_-$ is a zero of F iff $z \in i\mathbb{R}_-$ or $-z \in i\mathbb{R}_-$ is a zero of ξ.
vii) Let $g_n \neq \emptyset, n \geqslant 1$. The point $z \in g_n$ is a zero of F iff $z + i0 \in g_n^+$ or $z - i0 \in g_n^-$ is a zero of ξ (with the same multiplicity).
viii) Let $e_n^- > \max\{8e^{\|p\|_1}, 11C_*\}, C_* = \|q\|_t e^{\|p\|_1 + \|q\|_t + 2\|p\|_t}$ and $|g_n| > 0$. Then

$$F(e_n) \leqslant -\left(1 - \frac{R_1^2}{|e_n|^2}\right) \sinh^2 h_n < 0, \qquad \text{where} \qquad e_n = \sqrt{E_n} > 0, \tag{3.16}$$

and E_n, h_n are defined by (1.4). Moreover, F has at list two zeros in the segment $[e_n^-, e_n^+]$.

Proof. i) Using (3.10) we deduce that $F = \xi(z)\xi(-z)$ is entire and satisfies $F(z) = (1 - \Delta^2(z))(2 - A(z^2))^2 + J^2(z^2)$. Using (2.19) we obtain that S is entire and satisfies (3.15).

Recall that $F \geqslant 0$ and $S \geqslant 0$ inside each (e_{n-1}^+, e_n^-). Moreover, $F > 0$, since $a \neq 0$ inside each (e_{n-1}^+, e_n^-). Due to $F(e_n^\pm) \geqslant 0$, we get that F has even number of zeros on each interval $[e_n^-, e_n^+], n \geqslant 1$. We have $F = F_0 + S, F_0 = 4(1 - \Delta^2)$. Consider the case $z = e_n^+$, the proof for $z = e_n^-$ is similar. Thus if $F(z) = 0$, then we get $S(z) = 0$, since $\Delta^2(e_n^+) = 1$. Moreover, $F_0'(e_n^+) = -2\Delta(e_n^+)\Delta'(e_n^+) > 0$ and $S'(e_n^+) \geqslant 0$, which gives that $z = e_n^+$ is a simple zero of F.

ii) The function $m_\pm$ is analytic in $\mathcal{Z}_0$, then each $f^\pm(x, \cdot), x \in \mathbb{R}$ is analytic in $\mathcal{Z}_0$. Moreover, (2.7) yields $\beta(\mu_n) = \varphi(1, \mu_n) = \vartheta'(1, \mu_n) = 0$ and then $J(\mu_n^2) = 0$. Thus by (3.15), the function F has a double zero at μ_n at list.

iii) If $z \in g_n, n \geqslant 1$ is a bound state. Then (3.10) yields

$$0 = \xi(z + i0) = 2(-1)^{n+1} \sinh v(z + i0)(1 + A(z^2)) - J(z^2),$$

$$2 \sinh v(z + i0) = \frac{(-1)^{n+1} J(z^2)|}{(1 + A(z^2))} > 0.$$

If we assume that $z - i0$ is a anti-bound state, then $2 \sinh v(z - i0) = \frac{(-1)^{n+1} J(z^2)}{(1+A(z^2))} < 0$, which gives contradiction. The proof of iv) is similar.

v) The function ξ has not zeros in $\mathbb{C}_+ \setminus iR$, see [F1]. This yields v).

vi) The property $F(z) = F(-z), z \in \mathbb{C}$ gives vi).

vii) The statement vii) follows from iii).

viii) Estimate (3.12) gives $|A(e_n^2)| \leqslant \frac{1}{2}$ and (3.13) imply (3.16). The function $F(e_n^\pm) \geqslant 0$ and due to (3.16), we deduce that F has at list two zeros in the segment $[e_n^-, e_n^+]$. $\blacksquare$

Proof of Theorem 1.1. Identity (1.3) have been proved in Lemma 3.3.

i) The function ξ is analytic in $\mathcal{Z}_0$ and is real on $i\mathbb{R}$. Then the set of zeros of ξ is symmetric with respect to the imaginary line and satisfies (1.3).

The statement ii) has been proved in Lemma 3.3.

iii) Using Lemma 3.2, (2.26) and (2.27) we obtain

$$|A(z^2)| \leqslant \frac{\|q\|_t^2}{|z|^2} X_2, \qquad |J(z^2)| \leqslant \frac{3\|q\|_t}{|z|} X_2 X \qquad |\Delta(z) - \cos z| \leqslant \frac{X}{|z|}, \quad |z| \geqslant 2,$$

where $X_2 = e^{|\operatorname{Im} z|2t + \|q\|_t + 2\|p\|_t}$, $\quad X = e^{|\operatorname{Im} z| + \|p\|_1}$. Substituting these estimates into the identity

$$F - 4\sin^2 z = 4(\cos^2 z - \Delta^2) + J^2 + 4(1 - \Delta^2)A(A + 2)$$

we obtain

$$|F(z) - 4\sin^2 z| \leqslant 9X^2 C_0, \qquad C_0 = \frac{1}{|z|} + \frac{\|q\|_t^2 X_2^2}{|z|^2} + \frac{\|q\|_t^2 X_2}{|z|^2}\left(2 + \frac{\|q\|_t^2 X_2}{|z|^2}\right), \quad |z| \geqslant 2.$$

Using $e^{|\mathrm{Im} z|} \leqslant 4|\sin z|$ for all $|z - \pi n| \geqslant \frac{\pi}{4}, n \in \mathbb{Z}$, (see p. 27 [PT]), we obtain

$$9X^2 = 9e^{2|\operatorname{Im} z| + 2\|p\|_1} \leqslant |4\sin^2 z| r_0^2 \quad \textit{all } |z - \pi n| \geqslant \frac{\pi}{4}, \quad n \in \mathbb{Z}, \qquad r_0 = 6e^{\|p\|_1},$$

which yields

$$|F(z) - 4\sin^2 z| \leqslant 4|\sin^2 z| \frac{r_0^2 C_0}{|z|} < 4|\sin^2 z|, \quad \textit{all} \quad |z| \geqslant 2, z \notin \cup \mathcal{D}(\pi n, \frac{\pi}{4}),$$

since

$$\frac{r_0^2C_0}{|z|}\leqslant\frac{r_0^2}{|z|}+\frac{r_0^2\|q\|_t^2X_2^2}{|z|^2}+\frac{r_0^2\|q\|_t^2X_2}{|z|^2}\Big(2+\frac{\|q\|_t^2X_2}{|z|^2}\Big)<\frac{1}{5}+\frac{1}{4}+\frac{1}{2}+\frac{1}{(24)^2}<\frac{19}{20},$$

where we have used:

$$if\ \ z\in\mathcal{D}_F=\{z\in\mathbb{C}:|z|>\max\{180e^{2\|p\|_1},C_Fe^{2t|\operatorname{Im}z|}\}\}\quad\Rightarrow\quad\frac{r_0^2}{|z|}<\frac{1}{5},\quad\frac{r_0\|q\|_tX_2}{|z|}<\frac{1}{2},$$

and recall that

$$\mathscr{D}_F=\{\lambda\in\Lambda_2:|\lambda|^{\frac{1}{2}}>\max\{180e^{2\|p\|_1},C_Fe^{2t|\operatorname{Im}\sqrt{\lambda}|}\}\},\quad C_F=12\|q\|_te^{\|p\|_1+\|q\|_t+2\|p\|_t}.$$

Thus by Rouche's theorem, F has as many roots, counted with multiplicities, as $\sin^2 z$ in each disk $\mathcal{D}_{\frac{\pi}{4}}(\pi n)\subset\mathcal{D}_F$. Since $\sin z$ has only the roots $\pi n, n\geqslant 1$, then F has two zeros in each disk $\mathcal{D}_{\frac{\pi}{4}}(\pi n)\subset\mathcal{D}_F$ and F has not zeros in $\mathcal{D}_F\setminus\cup\mathcal{D}_{\frac{\pi}{4}}(\pi n)$.

$F(e_n^\pm)\geqslant 0$ for all n and Lemma 3.3 yields $F(e_n)<0$ for all n, where $e_n^->\max\{8e^{\|p\|_1},11C_*\}$, see (3.16). Then there exists exactly two simple real zeros $\varkappa_n^\pm=\sqrt{\lambda_n^\pm}>0$ of F such that $e_n^-\leqslant\varkappa_n^-<e_n<\varkappa_n^+\leqslant e_n^+$. Moreover, F has not no states in the "logarithmic" domain $\mathcal{D}_F\cap\mathbb{C}_-$.

The statements iv) and v) have been proved in Lemma 3.3. ■

4. PROOF OF THEOREM 1.2-1.4

Proof of Theorem 1.2. i) Theorem 1.1 gives $\varkappa_n^\pm=e_n^\pm\mp\delta_n^\pm=\sqrt{\lambda_n^\pm}$. Let $\varkappa=\varkappa_n^\pm,\delta=\delta_n^\pm$. Then the equation $0=\xi(\varkappa)=(-1)^{n+1}2(1+A(\varkappa^2))\sinh v(\varkappa)-J(\varkappa^2),\ \ \varkappa\in\overline{g_n^\pm}\neq\emptyset$ and (3.2) imply

$$\sinh|v(\varkappa)|=O(J(\varkappa^2))=\varepsilon O(|\varphi(1,\varkappa)|+\frac{|\vartheta'(1,\varkappa)|}{|\varkappa|^2}+\frac{|\beta(\varkappa)|}{\varkappa})=\varepsilon O(|g_n|),\quad\varepsilon=\frac{1}{2\pi n}$$

as $n\to\infty$. Moreover, using the estimate $|(z-e_n^-)(z-e_n^+)|^{\frac{1}{2}}\leqslant|v(z)|$ for each $z\in g_n$ (see [KK]) we obtain $|\delta(|g_n|-\delta)|^{\frac{1}{2}}\leqslant|v(\varkappa)|=\varepsilon O(|g_n|)$, which yields $\delta=\varepsilon^2O(|g_n|)$. Thus the points $\varkappa_n^\pm$ are close to $e_n^\pm$ and satisfy:

$$\delta_n^-=\varkappa_n^--e_n^-=\varepsilon^2O(|g_n|),\qquad and\qquad\delta_n^+=e_n^+-\varkappa_n^+=\varepsilon^2O(|g_n|).\tag{4.1}$$

Recall that $\varkappa_n^-$ and $\varkappa_n^+$ are simple. Consider the first case $\delta=\delta_n^-=|g_n|O(\varepsilon^2)$.

Using (3.11) we obtain

$$J=J_{10}+J_{10},\qquad J_{10}(z)=\int_{\mathbb{R}}\varphi(1,z,x)q(x)dx.$$

Let $\mu_n^2(\tau),\tau\in\mathbb{R}$ be the Dirichlet eigenvalue for the problem $-y''+q(x+\tau)y=z^2y,y(0)=y(1)=0$. In this case $\cos\phi_n$ in (1.7) is a function from $\tau\in\mathbb{R}$. Below we need facts from [K5]:

$$\frac{E_n^-+E_n^+}{2}-\mu_n^2(\tau)=\frac{|\gamma_n|}{2}\cos\phi_n(\tau),\quad\phi_n(\tau)=\phi_n(0)+2\pi n\tau+O(\varepsilon),\tag{4.2}$$

as $n\to\infty$ uniformly with respect to $\tau\in[0,1]$. Asymptotics (3.3) yield

$$\varphi(1,\varkappa,x)=\frac{(-1)^n}{\pi n}(1+O(\varepsilon))(\varkappa-\mu_n(x))=\frac{(-1)^n}{\pi n}(1+O(\varepsilon))(e_n^--\mu_n(x)+\varepsilon^2O(|g_n|)).\tag{4.3}$$

Thus we rewrite $e_n^- - \mu_n(x)$ in the form

$$e_n^- - \mu_n(x) = \frac{\frac{E_n^- + E_n^+ - |\gamma_n|}{2} - \mu_n^2(x)}{e_n^- + \mu_n(x)} = \frac{|\gamma_n|}{2} \frac{\cos\phi_n(x) - 1}{e_n^- + \mu_n(x)}.$$

This gives

$$J_{10}(\varkappa) = \int_{\mathbb{R}} \varphi(1,\varkappa,x)q(x)dx = \frac{(-1)^n}{\pi n}(1+O(\varepsilon)) \int_{\mathbb{R}} \left[\frac{|\gamma_n|}{2}\frac{\cos\phi_n(x)-1}{e_n^- + \mu_n(x)} + \varepsilon^2 O(|g_n|)\right] q(x)dx.$$

$$= \frac{(-1)^n |g_n|}{2\pi n} \int_{\mathbb{R}} \left[\cos y_n(x) - 1 + O(\varepsilon)\right] q(x)dx = \frac{(-1)^n|g_n|}{2\pi n}(-q_0 + c_n\widehat{q}_{cn} - s_n\widehat{q}_{sn} + O(\varepsilon)).$$

where $c_n = \cos y_n(0)$, $s_n = \sin y_n(0)$, and thus

$$(-1)^{n+1} J(\varkappa^2) = \frac{|g_n|}{2\pi n} I_n^-, \qquad I_n^- = q_0 - c_n\widehat{q}_{cn} + s_n\widehat{q}_{sn} + O(\varepsilon). \tag{4.4}$$

Using (3.12) we obtain

$$\sinh v(\varkappa) = \frac{(-1)^{n+1}J(\varkappa^2)}{2+2A(\varkappa^2)} = \frac{\varepsilon|g_n|I_n^-}{2+O(\varepsilon^2)}, \quad \operatorname{sign} v(\varkappa) = \operatorname{sign}(-1)^{n+1}J(\varkappa^2) = \operatorname{sign} I_n^-. \tag{4.5}$$

Note that if $v(\varkappa) > 0$ (or $v(\varkappa) < 0$), then $\varkappa \in g_n^+$ (or $\varkappa \in g_n^-$) and $\varkappa$ is a bound state (or a resonance). Moreover, if $v(\varkappa) = 0$, then $\varkappa = e_n^-$ or $\varkappa = e_n^+$ is a virtual state. Then (4.1) gives $\sinh v(\varkappa) = v(\varkappa)(1 + O(|g_n|^2\varepsilon^2))$ and using asymptotics (3.2), we obtain

$$v(\varkappa) = \sqrt{\delta(|g_n| - \delta)}(1 + O(\varepsilon^2)) = \sqrt{\delta|g_n|}(1+O(\varepsilon^2)).$$

Thus (4.5) yields $\delta_n^- = \frac{|g_n|\varepsilon^2}{4}(I_n^-)^2$ and (2.14) gives $|\gamma_n| = (2\pi n)|g_n|(1+O(\varepsilon^2))$, which yields $\delta_n^- = \frac{2|\gamma_n|}{(4\pi n)^3}(I_n^-)^2(1+O(\varepsilon^2))$.

If $q_0 > 0$, then $I_n^- > 0$ and above arguments yield that $\varkappa_n^-$ is a bound state and $\varkappa_n^+$ is an anti bound state. Conversely, if $q_0 < 0$, then $I_n^- < 0$ and we deduce that $\varkappa_n^-$ is an anti bound state and $\varkappa_n^+$ is a bound state.

Similar arguments imply the proof for the case $\delta_n^+ = e_n^+ - \varkappa_n^+ = |g_n|O(\varepsilon^2)$.

iii) We have $-c_n\widehat{q}_{cn} + s_n\widehat{q}_{sn} = -|\widehat{q}_n|\cos(\phi_n + \tau_n)$. Then (i) yields the statement (ii).

iv) An entire function $f(z)$ is said to be of *exponential type* if there is a constant α such that $|f(z)| \leqslant$ const. $e^{\alpha|z|}$ everywhere. The function f is said to belong to the Cartwright class $\mathscr{E}_\rho$, if f is entire, of exponential type, and the following conditions are fulfilled:

$$\int_{\mathbb{R}} \frac{\log^+|f(x)|dx}{1+x^2} < \infty, \quad \rho_\pm(f) = \rho, \quad \text{where} \quad \rho_\pm(f) \equiv \limsup_{y\to\infty} \frac{\log|f(\pm iy)|}{y}.$$

Denote by $\mathcal{N}^+(r,f)$ the number of zeros of f with real part $\geqslant 0$ having modulus $\leqslant r$, and by $\mathcal{N}^-(r,f)$ the number of its zeros with real part < 0 having modulus $\leqslant r$, each zero being counted according to its multiplicity. We recall the well known result (see [Koo]).

Theorem (Levinson). *Let the entire function $f \in \mathscr{E}_\rho, \rho > 0$. Then $\mathcal{N}^\pm(r,f) = \frac{r}{\pi}(\rho + o(1))$ as $r \to \infty$, and for each $\delta > 0$ the number of zeros of f with modulus $\leqslant r$ lying outside both of the two sectors $|\arg z|, |\arg z - \pi| < \delta$ is $o(r)$ for $r \to \infty$.*

Let $\mathcal{N}(r,f)$ be the total number of zeros of f with modulus $\leqslant r$. Denote by $\mathcal{N}_+(r,f)$ (or $\mathcal{N}_-(r,f)$) the number of zeros of f with imaginary part > 0 (or < 0) having modulus $\leqslant r$, each zero being counted according to its multiplicity.

Let $\pm\zeta_n > 0, n \in \mathbb{N}$ be all real zeros $\neq 0$ of F and let the zero $\zeta_0 = 0$ has the multiplicity $n_0 \leqslant 2$. Let $F_1 = z^{n_0} \lim_{r\to\infty} \prod_{|\zeta_n|\leqslant r}(1 - \frac{z}{\zeta_n})$. The Levinson Theorem and Lemma 2.3 imply

$$\mathcal{N}(r,F) = \mathcal{N}(r,F_1) + \mathcal{N}(r,F/F_1) = 2r\frac{1+2t+o(1)}{\pi}, \quad \mathcal{N}(r,F_1) = 2r\frac{1+o(1)}{\pi} \quad \text{as } r\to\infty.$$

Then Lemma 3.3 gives the identities $\mathcal{N}_-(r,F) = \mathcal{N}_+(r,F) = \mathcal{N}_-(r,\xi) + N_*$ for some integer $N_* \geqslant 0$. Thus we obtain

$$\mathcal{N}(r,F) = \mathcal{N}(r,F_1) + 2\mathcal{N}_-(r,\xi) + 2N_* = 2r\frac{1+2t+o(1)}{\pi}, \tag{4.6}$$

which yields $\mathcal{N}_-(r,\xi) = \frac{2rt+o(r)}{\pi}$ as $r\to\infty$ and (1.9).

Due to (1.8), the high energy states of H and H_0 are very close. This gives (1.10). ■

Proof of Theorem 1.3. i) Let the operator H_0 have infinitely many gaps $\gamma_n \neq \emptyset$ for some $p \in L^2(0,1)$ and let $\sigma = (\sigma_n)_1^\infty$ be any sequence, where $\sigma_n \in \{0,2\}$.

We take a potential q and let $\widehat{q}_n = \widehat{q}_{cn} + i\widehat{q}_{sn} = |\widehat{q}_n|e^{i\tau_n}, n \geqslant 1$ and $q_0 = 0$. We also assume that $|\widehat{q}_n| > n^{-\alpha}$ for n large enough and for some $\alpha \in (0,1)$. For each ϕ_n we take τ_n such that $|\cos(\phi_n + \tau_n)| > \varepsilon > 0$. Then due to Theorem 1.2 iii), the operator H has $\sigma_n = 1 - \operatorname{sign}\cos(\phi_n+\tau_n)$ bound states in the physical gap $\gamma_n^{(1)} \neq \emptyset$ and $2-\sigma_n$ resonances inside the nonphysical gap $\gamma_n^{(2)} \neq \emptyset$ for n large enough. Thus changing τ_n we obtain $\cos(\phi_n+\tau_n) < 0$, which yields $\sigma_n = 2$ or we obtain $\operatorname{sign}\cos(\phi_n + \tau_n) > 0$, which yields $\sigma_n = 0$.

ii) Let $q \in \mathcal{Q}_t, t > 0$ satisfy $q_0 = 0$ and let $|\widehat{q}_{cn}| > n^{-\alpha}$ for all n large enough and some $\alpha \in (0,1)$. Let $\sigma = (\sigma_n)_1^\infty$ be any sequence, where $\sigma_n \in \{0,2\}$. Let $\delta = (\delta_n)_1^\infty \in \ell^2$ be a sequence of nonnegative numbers $\delta_n \geqslant 0, n \geqslant 1$ and infinitely many $\delta_n > 0$.

Recall the result from [K5]:

The mapping $\Psi : \mathcal{H} \to \ell^2 \oplus \ell^2$ *given by* $\Psi = ((\Psi_{cn})_1^\infty, (\Psi_{sn})_1^\infty)$ *is a real analytic isomorphism between real Hilbert spaces* $\mathcal{H} = \{p \in L^2(0,1) : \int_0^1 p(x)dx = 0\}$ *and* $\ell^2 \oplus \ell^2$.

Then for the sequence $\delta = (\delta_n)_1^\infty \in \ell^2$ there exists a potential $p \in L^2(0,1)$ such that each gap length $|\gamma_n| = \delta_n$. Assume that $E_0^+ = 0$. Moreover, using Theorem 1.2 we deduce that H has exactly two simple states $\lambda_n^-, \lambda_n^+ \in \gamma_n^{(0)}$ for n large enough and λ_n^-, λ_n^+ have asymptotics

$$\sqrt{\lambda_n^\pm} = \sqrt{E_n^\pm} \mp \frac{2|\gamma_n|}{(4\pi n)^3}(R_n + O(1/n))^2, \qquad (-1)^{n+1}J(\lambda_n^\pm) = \frac{|\gamma_n|}{(2\pi n)^2}(R_n + O(1/n)),$$

where $R_n = -c_n\widehat{q}_{cn} + s_n\widehat{q}_{sn} = -|\widehat{q}_n|\cos(\phi_n + \tau_n)$ and $\widehat{q}_n = \widehat{q}_{cn} + i\widehat{q}_{sn} = |\widehat{q}_n|e^{i\tau_n}$ and ϕ_n is defined in (1.7). The parameters τ_n are fixed, but due to above results from [K5] the angles ϕ_n can be any numbers. If we take ϕ_n such that $|R_n| > \varepsilon > 0$ for all $\gamma_n \neq \emptyset$ and some $\varepsilon > 0$. Moreover, we take ϕ_n such that $\sigma_n = 1 - \operatorname{sign}\cos(\phi_n + \tau_n)$. Then Theorem 1.2 we obtain that the operator H has $\sigma_n = 1$ bound states in the physical gap $\gamma_n^{(1)} \neq \emptyset$ and $2 - \sigma_n$ resonances inside the nonphysical gap $\gamma_n^{(2)} \neq \emptyset$ for n large enough. ■

Acknowledgments. The various parts of this paper were written at ESI, Vienna, and Mathematical Institute of the Tsukuba Univ., Japan and Ecole Polytechnique, France. The author is grateful to the Institutes for the hospitality.

References

[BKW] B.M. Brown, I. Knowles, and R. Weikard, On the inverse resonance problem, J. London Math. Soc. (2) 68 (2003), no. 2, 383401.

[BW] Brown, B. M.; Weikard, R. The inverse resonance problem for perturbations of algebro-geometric potentials. Inverse Problems 20 (2004), no. 2, 481–494.

[Ch] Christiansen, T. Resonances for steplike potentials: forward and inverse results. Trans. Amer. Math. Soc. 358 (2006), no. 5, 2071–2089.

[F1] Firsova, N. Resonances of the perturbed Hill operator with exponentially decreasing extrinsic potential. Mat. Zametki 36 (1984), 711–724.

[F2] Firsova, N. The Levinson formula for a perturbed Hill operator. (Russian) Teoret. Mat. Fiz. 62 (1985), no. 2, 196–209.

[F3] Firsova, N. A direct and inverse scattering problem for a one-dimensional perturbed Hill operator. Mat. Sb. 130(172) (1986), no. 3, 349–385.

[Fr] Froese, R. Asymptotic distribution of resonances in one dimension, J. Diff. Eq., 137(1997), 251-272.

[GT] Garnett, J.; Trubowitz, E. Gaps and bands of one dimensional periodic Schrödinger operators II. Comment. Math. Helv. 62(1987), 18-37.

[GS] Gesztesy, F.; Simon, B. A short proof of Zheludev's theorem. Trans. Amer. Math. Soc. 335 (1993), no 1, 329–340.

[H] Hitrik, M. Bounds on scattering poles in one dimension. Comm. Math. Phys. 208 (1999), no. 2, 381–411.

[KK] Kargaev, P.; Korotyaev, E. Effective masses and conformal mappings. Comm. Math. Phys. 169 (1995), no. 3, 597–625.

[KK1] Kargaev, P.; Korotyaev, E. Inverse Problem for the Hill Operator, the Direct Approach. Invent. Math., 129(1997), no. 3, 567-593.

[KM] Klopp, F.; Marx, M. The width of resonances for slowly varying perturbations of one-dimensional periodic Schrödingers operators, Seminaire: Equations aux Derivees Partielles. 2005–2006, Exp. No. IV, 18 pp., Semin. Equ. Deriv. Partielles, Ecole Polytech., Palaiseau.

[K1] Korotyaev, E. Inverse resonance scattering on the half line, Asymptotic Anal. 37(2004), No 3/4, 215-226.

[K2] Korotyaev, E. Inverse resonance scattering on the real line. Inverse Problems 21 (2005), no. 1, 325–341.

[K3] E. Korotyaev, Stability for inverse resonance problem. Int. Math. Res. Not. 2004, no. 73, 3927–3936.

[K4] Korotyaev, E. Resonances for Schrödinger operator with periodic plus compactly supported potentials on the half-line, preprint 2008.

[K5] Korotyaev, E. Inverse problem and the trace formula for the Hill operator. II Math. Z. 231 (1999), no. 2, 345–368.

[Koo] Koosis, P. The logarithmic integral I, Cambridge Univ. Press, Cambridge, London, New York 1988.

[M] Marchenko, V. Sturm-Liouville operator and applications. Basel: Birkhäuser 1986.

[PT] Pöschel, P.; Trubowitz, E. Inverse Spectral Theory. Boston: Academic Press, 1987.

[Rb] Rofe-Beketov, F. S. A perturbation of a Hill's operator, that has a first moment and a non-zero integral, introduces a single discrete level into each of the distant spectral lacunae. (Russian) Math. Phys. i Func. Analiz. Kharkov, No 4, 1973, 158–159, 163.

[S] Simon, B. Resonances in one dimension and Fredholm determinants. J. Funct. Anal. 178(2000), 396–420.

[So] Sobolev, A.V. Weyl asymptotics for the discrete spectrum of the perturbed Hill operator. Estimates and asymptotics for discrete spectra of integral and differential equations (Leningrad, 1989–90), 159–178, Adv. Soviet Math., 7, Amer. Math. Soc., Providence, RI, 1991.

[T] Titchmarsh, E. Eigenfunction expansions associated with second-order differential equations 2, Clarendon Press, Oxford, 1958.

[Tr] Trubowitz, E. The inverse problem for periodic potentials. Commun. Pure Appl. Math. 30(1977), 321-337.

[Z] Zworski, M. Distribution of poles for scattering on the real line, J. Funct. Anal. 73(1987), 277-296.

[Z1] Zworski, M. SIAM, J. Math. Analysis, "A remark on isopolar potentials" 82(6), 2002, 1823-1826.

[Z2] Zworski, M. Counting scattering poles. In: Spectral and scattering theory (Sanda, 1992), 301331, Lecture Notes in Pure and Appl. Math., 161, Dekker, New York, 1994.

[Zh] Zheludev, V. A. The eigenvalues of a perturbed Schrdinger operator with periodic potential. (Russian) 1967 Problems of Mathematical Physics, No. 2, Spectral Theory, Diffraction Problems pp. 108–123.

School of Math., Cardiff University. Senghennydd Road, CF24 4AG Cardiff, Wales, UK. email KorotyaevE@cf.ac.uk, Partially supported by EPSRC grant EP/D054621.